



**POWERCUBES™ 28% + RUM  
TYPE C MEDICATED FEED**

**PROTEIN SUPPLEMENT  
for MATURE REPRODUCING BEEF COWS  
(on Pasture or Dry Lot)**

**For improved feed efficiency when receiving supplemental feed; For prevention and control of coccidiosis caused by Eimeria bovis and Eimeria zuernii.**

**ACTIVE DRUG INGREDIENT**

Monensin (as Monensin sodium) 0.000165% or 150 g/ton  
Each pound contains 75 mg Monensin

**TYPICAL ANALYSIS**

Crude Protein	(Min)	28.0 %
Crude Fat	(Min)	6.0 %
Crude Fiber	(Max)	10.0 %
Acid Detergent Fiber	(Max)	18.0 %
Calcium	(Min)	0.7%
Calcium	(Max)	1.0%
Phosphorus	(Min)	0.6 %
Selenium	(Min)	0.3 ppm

**FEEDING DIRECTIONS**

**For coccidiosis prevention and control:** Hand feed at the rate of 0.14 to 0.42 mg monensin/lb/BW per day, depending upon severity of challenge, up to 200 mg monensin per head per day.

**For improved feed efficiency:** Hand feed or mix 0.67 lb to 2.67 lb of POWERCUBES™ + RUM per head per day (to provide 50 to 200 mg monensin per head per day) into the total ration and feed continuously. Feed undiluted or blend with an appropriate amount of ground feed. Feed (other than POWERCUBES™ + RUM) can be restricted to 95% (of normal requirements) when 50 mg of monensin is fed, and to 90% at 200 mg monensin.

Cows on pasture or in dry lot are required to receive their daily drug amount in a minimum of 1 lb of Type C Medicated Feed. Additionally, a minimum of 16 lb (air-dry basis) of roughage such as silage, haylage, ammoniated straw, hay or equivalent feedstuffs should be fed in order to meet NRC recommendations for mature reproducing beef cows to gain 0.25 to 0.75 lb per head per day.

Standing, dried winter range forage may not be of adequate quality to result in improved efficiency when supplemented with monensin. In all cases, provide plenty of fresh, clean water, adequate forage, and a source of vitamins and minerals. For further feeding directions, consult your supplier or nutritionist.

**FEEDING TABLE**

Recommended POWERCUBES™ + RUM (75 mg/lb monensin) Feeding Level (lb/head/day)	Monensin Amount (mg/head/day)
0.67	50
1	60
2.67	200**

\*\*Maximum Approved Level

**IMPORTANT**

During the first 5 days, cattle should receive no more than 100 mg monensin/day contained in not less than 1 lb of feed.

**CAUTION**

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

**WARNING**

A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.



NADA#095-735